Prunus avium in Europe: distribution, habitat, usage and threats

E. Welk, D. de Rigo, G. Caudullo

Prunus avium (L.) L., known as wild cherry, is a medium sized, fast growing and rather short-lived deciduous tree. The wild populations, described here, belong to the typical subspecies avium, whereas the cultivated forms are distinguished as subsp. duracina and subsp. juliana. The early large white flowers are clustered on spur shoots and give rise to edible purplish small drupes with a bitter-sweet taste. The main stem trunk is usually very straight with a characteristic greyreddish-brown "cherry-bark" that is shiny with large horizontal lenticels and horizontally peeling pattern. Wild cherry occurs as a minor component in many types of temperate broadleaved and mixed forests. The wild form is a mainly European species and served as the origin of all cultivated forms. It is a very popular ornamental fruit tree, and the hard, reddish-brown timber is one of the most valuable in Europe.

Wild cherry (Prunus avium (L.) L.) is a fast-growing but short-lived (100-150 years), medium sized deciduous tree, which grows to 15-32 m height and with a stem diameter of up to 90-120 cm¹. The species mostly develops single, straight trunks with a thin, smooth purplish-grey bark that becomes grey-brown with horizontal fissuring and peeling when old. Young trees grow with a strong apical control developing a straight trunk and an erect-pyramidal "coniferous" crown shape, becoming broader and rounded on single old trees or conical in individuals in forest stands^{2, 3}. Young shoots are shiny, pale grey to purplish-brown, and have large, reddish brown and protruding ovoid-ellipsoid, glabrous winterbuds at the branch ends arranged in whorllike form. Stem wounds produce a resin-like, amber coloured odourless gum^{4, 5}. The leaves change colour from light green in spring over dark green in summer, and to yellow, orange-red, scarlet or pink in autumn. They are alternate, pendulous, simple and elliptic-ovate to obovate acute in shape. The leaf margins are mildly serrated with slightly rounded teeth. There are conspicuous pairs of dark-red glands at the 2-3.5 cm long petiole below the lamina. Leaf size is approximately 5-15x3-8 cm. They are usually dull, glabrous-rugose above and sometimes weakly downy at the 8-15 pairs of secondary vein ribs beneath^{1, 2, 5}. Wild cherry flowers are allogamous, actinomorphic, about 2-2.5 cm in diameter, white, hermaphroditic, insect pollinated, and are arranged in **racemose** clusters of 2-5 flowers on short spurs (brachyblasts) with multiple apical (inserted at tips) buds; of which the distal (uppermost) bud is vegetative and continues growth, while the others bear new inflorescences³. Flowers are pollinated mainly by honeybees, wild bees and bumblebees and the trees are generally not self-fertile^{1, 2, 5}. Individual trees have a relatively short life span of 100-120 years at maximum, and can start fruiting when 10-15 years old 6. In Central Europe, flowering starts earliest in late March and occurs until May, while



Frequency of Prunus avium occurrences within the field observations as reported by the National Forest Inventories. The chorology of the native spatial range for *P. aviumis* derived after EUFORGEN¹⁹.

individual trees are in flower for about one week^{1, 2, 5}. Fruits are purplish black drupes, sub-globose to ovoid, 1-2 cm in diameter with a smooth, fleshy, and bitter-sweet edible endocarp^{1, 2, 5}. Ripe fruits occur from late spring until summer and are consumed and dispersed mostly by birds such as pigeons, starlings, thrushes and jays, but also by larger vertebrates like foxes, badgers or wild boar^{2, 6}.





.... Cutivated cherry tree in orchard. (Copyright Tara2, commons.wikimedia.org: CC-BY)

Distribution

Prunus avium occurs naturally throughout the temperate forest regions of Europe, Anatolia, and adjacent regions of the North African Maghreb, and western Asia^{1, 2, 5}. The distribution area extends northwards to the British Islands and to southern Scandinavia, where it is difficult to tell apart native from naturalised populations. The northern natural range limit is reached at about 55° north parallel¹. In the south the range extends to North Africa, South Spain, central Italy, and the Balkans. In these regions it is strictly confined to increasingly scattered high altitude populations in humid situations^{1, 2}. The easternmost parts of the natural range include the Caucasus and the Elburs mountains of North Iran, where the species is reported to occur up to elevations of 2000m. In the Alps it occurs up to 1700 m and reaches 1900 m in South-East France⁷. The general altitudinal distribution stretches from the planar to submontane altitudinal zone. In the highest, montane locations wild cherry often grows only as a shrub⁷. Limiting factors for wild cherry distribution are mainly related to rainfall in the summer period in the south and colder conditions in north and east Europe⁸. Beside the circumscribed native distribution, this Prunus is widely planted and naturalises successfully in deciduous forest habitats and shrub land, especially in temperate regions of Northern Asia and North America^{1, 2}.



..... Map 2: High resolution distribution map estimating the relative probability of presence

Reddish-brown bark with large horizontal lenticels in stripes (Convright Stefano Zerauschek w

Habitat and Ecology

Wild cherry is a mesophytic, comparatively shallow-rooting, light demanding species, which can grow in quite different soil types. However, it favours deep fertile soils with a good water supply. The tree does not tolerate heavy clays, waterlogged or poorly drained sites and can be sensitive to drought^{1, 2, 5}. Main habitat type is semi-shade, open deciduous woodland or scrubland especially at edges, glades and clearings, where this tree essentially occurs as a rare and scattered pioneer species^{1, 2}. The pioneer colonisation strategy is realized as a first generation establishment via seedling recruitment, potentially followed by sometimes extensive vegetative growth via root suckering. With its ability for coppice

Prunus avium

shooting and sucker formation as well as its rapid juvenile growth, wild cherry possesses competitive advantages in early succession stages. In natural forest stands the species is usually replaced by climax tree species during ongoing succession^{9, 10}. In its European main distribution range it is a frequent element of several mixed deciduous forests type alliances of the class Querco-Fagetea, such as ravine forests (Tilio-Acerion), oak-Hornbeam forests (Carpinion betuli), lowland beech forests (Fagion), and riverine floodplain forests (Alno-Ulmion)¹¹.



.... Flowers are arranged in clusters inserted at tips of buds (Copyright AnRo0002, commons.wikimedia.org: CC0)

Importance and Usage

Wild cherry fruits have been a source of food for humans for several thousands of years^{1, 5}. Today it is cultivated as a fruit tree in temperate regions all over the world. Beside human fruit consumption in the cultivated subspecies, wild cherry is also one of the most important European hardwood trees, with a valuable solid and dense wood that is highly sought after for panelling and cabinet-making, and also suitable for producing parquet floors and musical instruments. Today, veneer production is the main use of wild cherry timber⁶. Beside its usage as an ornamental landscape tree, this tree is also an important food source for many species of songbirds and insects^{2, 5}. The distribution range of wild cherry, and even its suitable potential range for silvoarable agroforestry¹², overlaps with many areas in Europe with high erosion rates such as the European mountain systems¹³. Its adventitious roots are suitable to be exploited for deep reinforcement and soil strength enhancement¹⁴ as well as for soil bioengineering to increase the stability of slopes and mitigate erosion¹⁵.

Threats and Diseases

Prunus avium develops a rather shallow heart-shaped root system with far reaching lateral roots in top soil horizons, rendering it quite vulnerable to wind-throw. Additionally it is relatively sensitive to environmental stresses and in unfavourable conditions, so that it can easily be attacked by a variety of pests and diseases^{1, 2, 5}. Roots may be attacked by mice and voles, and leaves by caterpillars of e.g. winter moth (Operophtera brumata), while the larvae of European cherry fruit fly (Rhagoletis cerasi) and the bird-cherry weevil (Anthonomus rectirostris) feed on the fruits^{2, 5}. As most species of the genus *Prunus*, wild cherry is vulnerable to the gypsy moth (*Lymantria dispar*)^{16, 17}. Infectious diseases may be cherry leaf roll virus (CLRV), bacterial cankers like Pseudomonas syringae or fireblight (Erwinia amylovora)¹⁸. Common foliar fungal pathogens are leaf scorch (Apiognomonia *erythrostoma*) and leaf spot (*Blumeriella jaapi*)¹. Young wild cherry trees are especially susceptible to browsing by ungulate herbivores^{2, 5}.

Field data in Europe (including absences)
Observed presences in Europe





Map 3: High resolution map estimating the maximum habitat suitability

References

- A. Praciak, et al., The CABI encyclopedia of forest trees (CABI, Oxfordshire, UK, 2013). [1] [2] T. Schmid. Enzyklopädie der
- Holzgewächse: Handbuch und Atlas der Dendrologie, A. Roloff, H. Weisgerber, U. M. Lang, B. Stimm, P. Schütt, eds. (Wiley-Vch Verlag, Weinheim, 2006).
- D. Barthélémy, Y. Caraglio, S. Sabatier, [3] Valuable Broadleaved Forests in Europe, H. Spiecker, S. Hein, K. Makkonen-Spiecker, M. Thies, eds. (Brill Academic Publishers, Leiden, 2009), pp. 87-101
- A. Kurtto, Euro+Med Plantbase [4] the information resource for Euro Mediterranean plant diversity (2009) http://www.emplantbase.org.
- H. Scholz, I. Scholz, Gustav Hegi [5] Illustrierte Flora von Mitteleuropa, Band 4 Teil 2B: Rosaceae, H. Scholz, ed. (Blackwell senschafts-Verlag, Berlin, 1995), pp 446–510, second edn.
- F. Ducci, B. De Cuyper, A. De Rogatis, J. [6] Dufour, F. Santi, *Forest Tree Breeding in Europe*, L. E. Pâques, ed. (Springer Netherlands, 2013), vol. 25 of Managing Forest Ecosystems, pp. 463–511.
- P. Schwab, Kirschbaum Projekt Förderung seltener Baumarten (Professur Waldbau of Phytopathology **27**, 165 (1989). ETHZ, 2001).
- K. Russell, EUFORGEN Technical Guidelines for aenetic conservation and use for wild cherry (Prunus avium) (Bioversity International, 2003).

- [9] S. P. Vaughan, J. E. Cottrell, D. J. Moodley, T. Connolly, K. Russell, *Forest Ecology and Management* 242, 419 (2007).
- [10] A. M. Höltken, H.-R. Gregorius, BMC Ecology 6, 1 (2006).
- [11] W. Willner, *Phytocoenologia* **3**, 337 (2002).
- [12] Y. Reisner, R. de Filippi, F. Herzog, J. Palma, Ecological Engineering 29, 401 (2007). [13] C. Bosco, D. de Rigo, O. Dewitte, J. Poesen,
- P. Panagos, Natural Hazards and Earth System Science **15**, 225 (2015). [14] J. E. Norris, A. Di Iorio, A. Stokes, B. C.
- Nicoll, A. Achim, Slope Stability and Erosion Control: *Ecotechnological Solutions*, J. E. Norris, *et al.*, eds. (Springer Netherlands, 2008), pp. 167–210.
- [15] F. Florineth, H. P. Rauch, H. Staffler, Proceedings of the International Congress INTERPRAEVENT 2002 in the Pacific Rim (2002), vol. 2, pp. 827-837.
- [16] D. de Rigo, et al., Scientific Topics Focus 2, mri10a15+ (2016).
- [17] H. Kutinkova, R. Andreev, *Journal of Fruit* and Ornamental Plant Research **12**, 41 (2004).
- [19] EUFORGEN, Distribution map of wild
- cherry (Prunus avium) (2008) www.euforgen.org

..... White flowers are insect pollinated. (Copyright mornarsamotarsky, www.flickr.com: CC-BY)

·..· Autoecology diagrams based on harmonised field observations from forest plots





Cherres are ripe from late spring until summer and are dispersed principally by birds. (Copyright Steven Gill, www.flickr.com: CC-BY)

This is an extended summary of the chapter. The full version of this chapter (revised and peer-reviewed) will be published online at https://w3id.org/mtv/FISE-Comm/v01/e01491d. The purpose of this summary is to provide an accessible dissemination of the related main topics.

This QR code points to the full online version, where the most updated content may be freely accessed.

Please, cite as:

Welk, E., de Rigo, D., Caudullo, G., 2016. Prunus avium in Europe distribution, habitat, usage and threats. In: San-Miguel-Ayanz, J., de Rigo, D., Caudullo, G., Houston Durrant, T., Mauri, A. (Eds.), Europe an Atlas of Forest Tree Species. Publ. Off. EU, Luxembourg, pp. e01491d+

