Prunus spinosa in Europe: distribution, habitat, usage and threats

I. Popescu, G. Caudullo

The blackthorn (*Prunus spinosa* L.) is a spiny, deciduous shrub which produces small, purple, edible plums. This species occurs mostly from south-central Europe up to southern Scandinavia, and eastwards to Asia Minor, growing in forest margins and open woodlands as part of Mediterranean thermophilous plant communities. It is cultivated as an ornamental plant and for fruit production, used to make jams, wine, vinegar and distillates. The blackthorn has no important threats, but it can be a natural host and potential reservoir of diseases affecting production of economically important fruits, such as apricots, plums, peaches and apples.

The blackthorn, or sloe, (Prunus spinosa L.) is a spiny, deciduous shrub, growing 1-5 m tall. It forms a dense canopy with intricate branches and numerous suckers¹⁻⁴. Secondary twigs often transformed into a spine, initially velvety soft, reddishbrown. The buds are globular oval, reddish-brown, more or less hairy. The bark is dark grey to blackish, slightly grooved. The leaves are alternate, 2-5 × 1-2 cm long, obovate to oblanceolate, or elliptical, with margins finely toothed, dull green in colour and hairless above, usually hairy on the veins underneath^{1, 3}. The petioles are 0.2-1 cm long, often hairy. The stipules are elongate, glandular, toothed, and usually longer than petioles³. The flowers are white, 1-1.7 cm wide, usually solitary, appearing before leaves, numerous, on about 0.5 cm long pedicels¹⁻³. The sepals are triangular-ovate, often glandular toothed. Petals have a short claw (a thinner part at the base of the petal). The stamens are usually 20, about 0.5 cm long. The anthers are yellow or red³. The fruit is 1-1.5 cm drupe, globose, purple covered with a frostlike bloom, ripening bluish-black, pulp greenish, sour and astringent, not easily detaching from the endocarp. This species flowers in March-May. The fruits ripen in late summer and autumn, and are sometimes persistent on the plant through winter^{1-3, 5}.

Distribution

This species occurs in most of South-Central Europe, except the lower half of the Iberian Peninsula, extending northwards to the southern part of the Scandinavian Peninsula. Eastwards it reaches Asia Minor, Caucasus and the Caspian Sea. There are also isolated populations in Tunisia and Algeria^{1, 6}. It has been introduced and locally naturalised in North America⁷ and New Zealand⁸. It grows from lowlands to 1600m on Southern Alps (Switzerland)^{6, 9}.





 Frequency of *Prunus spinosa* occurrences within the field observations as reported by the National Forest Inventories. The chorology of the native spatial range for *P. spinosa* is derived after several sources^{6, 27-31}.

Habitat and Ecology

The blackthorn occurs in forest margins and openings, on sunny, rocky slopes, ravines, river valleys, in meadows and pastures from low elevations to the mountains^{1, 2, 10}. It is one of the main species of scrub, a belt of shrubs adjacent to the forest, in the ecotone between woodland and grassland communities^{11, 12}. These Euro-Siberian and Mediterranean thermophilous communities (class Rhamno-Prunetea) are dominated by thorny and spiny, scrambling shrubs, developing in mesic to dry soils at the edges of oak and beech forests or on river banks with willows and poplars. Rarely are scrubs natural and permanent, more commonly they are secondary vegetation developing on disturbed sites or stages of secondary succession in abandoned meadows or pastures at the borders of the agricultural landscape. The blackthorn is commonly found with other shrub species of the genera Rosa, Rubus, Prunus and Cornus, as well as with spindle tree (*Euonymus europaeus*), hawthorn (*Crataegus* monogyna), wild privet (Ligustrum vulgare), etc.^{10, 13-16}

Importance and Usage

As with most Prunus species, the blackthorn has a high level of hybridisation and crossed with European plum (Prunus *domestica*) it forms the hybrid *Prunus* x *fruticans*. This species is considered one of the progenitors of the cultigen European plum together with cherry plum (*Prunus cerasifera*)¹⁷. Fully ripe fruits even though astringent are edible, and mainly used to make jams, jellies, preserves, wine, vinegar, and distilled alcoholic beverages, or as ingredients of various pastries. Flowers and petals in particular are used medicinally as tea, syrup, fresh juice or tincture to fight diarrhoea, anaemia, and other ailments^{3, 5}. Due to their vitamin C and the phenolic acids content, the fruits can be a valuable source of natural antioxidants¹⁸. The blackthorn is used as an ornamental tree, often as hedges^{3, 9, 10, 13}. It is also an important plant for wildlife, its early spring flowers provide nectar for early pollinators, its branches create a spiny thicket, providing secure nesting for birds and protection and food for small mammals^{13, 19, 20}. In temperate oak forests, its dense and thorny canopy is difficult to penetrate by large herbivores. Together with other scrub species, thorny shrubs can protect young palatable seedlings, such as oak seedlings, against browsing along the forest edges and clearings^{21, 22}.



 Purple globose drupes covered with a frostlike bloom. (Copyright Phil Sellens, www.flickr.com: CC-BY)



 Hermaphrodite flowers appear in spring and have 5 white petals with 20 stamens.
 (Copyright Maia Dumat, www.flickr.com; CC-BY)

References

- D. A. Webb, Flora Europea. Volume 2. Rosaceae to Umbelliferae, T. G. Tutin, et al., eds. (Cambridge University Press, 1968), pp. 77–80.
- [2] O. Polunin, *Flowers of Europe: A Field Guide* (Oxford University Press, 1969).
- [3] T. Săvulescu, ed., Flora Republicii Populare Romil,ne, vol 4 (Editura Academiei Române, Bucureşti, 1956).
- [4] O. Johnson, D. More, *Collins tree guide* (Collins, 2006).
- [5] B.-E. van Wyk, Food Plants of the World: An Illustrated Guide (Timber Press, 2005)
 [6] H. Meusel, E. Jager, S. Rauschert, E.
- 6] H. Meusel, E. Jager, S. Rauschert, E. Weinert, Vergleichende Chorologie der Zentraleuropäischen Flora (Gustav Fischer Verlag Jena, 1978).
- [7] USDA NRCS, The PLANTS database (2015). National Plant Data Team, Greensboro, USA, http://plants.usda.gov.
- [8] P. B. Heenan, et al., New Zealand Journal of Botany **37**, 629 (1999).
- [9] V. L. Komarov, et al., Flora of the USSR - Volume X: Rosaceae - Rosoideae, Prunoideae, vol. 10 of Flora of the USSR
- (Keter Press, Jerusalem, 1970). [10] H. Ružičková, L. Halada, L. Jedlička, E.

- [16] U. Bohn, et al., Karte der natürlichen Vegetation Europas; Map of the
- Natural Vegetation of Europe (Landwirtschaftsverlag, 2000). [17] M. B. Crane, W. J. Lawrence, *The Genetics* of Garden Plants (MacMillan & Company, London, 1952), fourth edn.
 - [18] B. Marìa Ruiz-Rodrìguez, *et al., Fruits* **69**, 61 (2014).
 - [19] P. Freschi, et al., European Journal of Wildlife Research 60, 423 (2014).
 - [20] P. Freschi, et al., Mammalia **79**, 51 (2015).
 [21] F. W. M. Vera, *Grazing Ecology and Forest History* (CABI, 2000).
 - [22] E. S. Bakker, et al., Journal of Applied Ecology 41, 571 (2004).
 - [23] P. Salamon, L. Palkovics, European Journal of Plant Pathology 108, 903 (2002).
 [24] M. G. Podrigues, A. Eonseca, International.
 - [24] M. G. Rodrigues, A. Fonseca, International Journal of Systematic and Evolutionary Microbiology 53, 607 (2003).
 - [25] H. İlbağı, A. Çıtır, H. Bostan, Acta Horticulturae **781**, 33 (2008).
 - [26] M. Rankovic, I. Dulic-Markovic, Acta Horticulturae **309**, 151 (1992).
 [27] A. A. Dönmez, c. Yildirimli, *Turkish Journal*
 - of Botany **24**, 187 (2000).

••• Spiny branches with fruits in autumn. (Copyright Giovanni Caudullo: CC-BY)

Threats and Diseases

Despite being a member of the *Prunus* genus, the blackthorn has no important threats. It can be a natural host and potential reservoir of diseases affecting production of economically important fruits, such as apricots, plums, peaches and apples, and tests have revealed that the presence of wild blackthorns nearby could be the cause of increased presence of fruit viruses (e.g. Sharka, dwarf virus, necrotic ring spot virus, chlorotic leaf spot virus) and fungi (e.g. *Taphrina pruni*)²³⁻²⁶.

krajinnej ekológie SAV, Bratislava, 1996).

- [11] J. G. Kelcey, N. Müller, eds., Plants and Habitats of European Cities (Springer-Verlag, New York, 2011).
- [12] S. R. Mortimer, et al., The nature conservation value of scrub in Britain Tech. rep., Joint Nature Conservation Committee (2000). N. 308.
- [13] I. Žgančiková, T. Vereš, T. Baranec, Research Journal of Agricultural Sciences 44, 233 (2012).
- [14] J. C. Costa, *et al., Global Geobotany* **2**, 1 (2012).
- [15] M. Chytrý, ed., Vegetace České republiky 4: Lesnì a křovinná vegetace - Vegetation of the Czech Republic 4: Forest and scrub vegetation (Academia, Praha, 2013).
- -____
- [28] A. N. Afonin, S. L. Greene, N. I. Dzyubenko, A. N. Frolov, eds., Interactive Agricultural Ecological Atlas of *Russia and Neighboring Countries: Economic Plants and their Diseases, Pests and Weeds* [Online] (2008). http://www.agroatlas.ru.
- [29] Sociedade Portuguesa de Botânica, Flora-On: Flora de portugal interactiva (2014). http://www.flora-on.pt.
- [30] Anthos, Information System of the plants of Spain (Real Jardin Botánico, CSIC -Fundación Biodiversidad, 2015). http://www.anthos.es.
- [31] Botanical Society of Britain & Ireland, *BSBI Big Database* (2015). http://bsbidb.org.uk.

This is an extended summary of the chapter. The full version of this chapter (revised and peer-reviewed) will be published online at https://w3id.org/mtv/FISE-Comm/v01/e018f4e. The purpose of this summary is to provide an accessible dissemination of the related main topics.

This QR code points to the full online version, where the most updated content may be freely accessed.

Please, cite as

Popescu, I., Caudullo, G., 2016. *Prunus spinosa* in Europe: distribution, habitat, usage and threats. In: San-Miguel-Ayanz, J., de Rigo, D., Caudullo, G., Houston Durrant, T., Mauri, A. (Eds.), *European Atlas of Forest Tree Species*. Publ. Off. EU, Luxembourg, pp. e018f4e+

